

A two-dimensional model for the dynamics of sand patches

Camille Rambert¹, Clément Narteau², Joanna Nield³, Giles Wiggs⁴, Pauline Delorme⁵, Matthew Baddock⁶, Philippe Claudin⁷

¹ Physique et Mécanique des Milieux Hétérogènes, ESPCI Paris, Paris, France

² Institut de physique du Globe de Paris, Université de Paris, CNRS, Paris, France

³ School of Geography and Environmental Science, University of Southampton, Southampton, UK

⁴ School of Geography and the Environment, University of Oxford, Oxford, UK

⁵ Energy and Environment Institute, University of Hull, Hull, UK

⁶ Geography and Environment, Loughborough University of Technology, Loughborough, UK

⁷ Physique et Mécanique des Milieux Hétérogènes, CNRS, ESPCI Paris, PSL Research University, Sorbonne Université, Université de Paris, Paris, France

`camille.rambert@espci.fr`

Sand patches are one of the early stages of aeolian bedforms. They form on non-erodible surfaces in both desert and coastal environments. Their initiation is associated with the change of saltation transport law on rigid and granular beds [1]. Here we present a two-dimensional model that couples these surface-dependent transport laws with the feedback of the bed elevation on the wind flow. Analysing the spatio-temporal evolution of an initial very flat sand patch, we emphasise the central role of the input flux as well as the lengthscale over which occurs the transition between the two transport laws. We also show that, for adjusted parameters of the model, we are able to reproduce the growth and the propagation of these small metre-scale bedforms over time, in quantitative comparison with field measurements.

Références

1. P. DELORME, J.M. NIELD, G.F.S. WIGGS, M.C. BADDOCK, N.R. BRISTOW, J.L. BEST, K.T. CHRISTENSEN, P. CLAUDIN, *Geophysical Research Letters*, **50** (2023).